

DOI	10.12691/ajbr-4-4-3
Title	Drug resistant <i>Staphylococcus aureus</i> in clinical samples at Kampala International University-teaching hospital, Bushenyi District, Uganda.
Authors	Janet Nalwoga, Michael Tirwomwe, Albert Nyanchoka Onchweri, Josephat Nyabayo Maniga, Cyprian Mose Nyaribo, Conrad Ondieki Miruka
Journal	<i>American Journal of Biomedical Research</i>
Year	2016
Abstract	Contexte : <i>Staphylococcus aureus</i> résistant à la méthicilline (SARM) est un pathogène nosocomial majeur qui provoque souvent des infections difficiles à traiter. Cela est dû au fait que ce pathogène est généralement résistant à d'autres antibiotiques couramment utilisés. La présence de SARM pharmacorésistant chez les patients a déjà été documentée dans diverses régions de l'Ouganda. Cependant, aucun rapport n'a été enregistré pour l'hôpital universitaire de l'Université Internationale de Kampala (<i>Kampala International University Teaching Hospital</i>) dans l'ouest de l'Ouganda. Cette étude a donc été menée pour déterminer la prévalence et la résistance aux antibiotiques des souches de SARM chez les patients hospitalisés dans le service de chirurgie. Méthodes : Des écouvillons de plaies ont été prélevés sur des patients des deux sexes hospitalisés dans le service de chirurgie. Les échantillons ont ensuite été mis en culture sur des milieux appropriés. Les colonies de <i>Staphylococcus aureus</i> obtenues ont été testées pour leur résistance à l'oxacilline afin d'identifier les souches de SARM. De plus, la résistance aux antibiotiques des isolats de SARM a été déterminée par la méthode de diffusion par disque en utilisant divers antibiotiques. Résultats : Sur les cent quatorze isolats issus des échantillons cliniques, seuls soixante-quinze isolats ont été clairement identifiés comme <i>S. aureus</i>, dont 85,3 % de coagulase positive et 13,3 % de coagulase négative. La prévalence de <i>Staphylococcus aureus</i> résistant à la méthicilline était de 56,1 %. Parmi les isolats de SARM, la résistance à la ciprofloxacine s'est avérée être la plus élevée, tandis que la résistance à la ceftriaxone s'est avérée être la plus faible. Conclusion : La prévalence élevée de SARM chez les patients du service de chirurgie nécessite la prise de mesures appropriées pour empêcher une propagation plus large du pathogène. Il est recommandé de déterminer la source de ces souches de SARM pharmacorésistantes afin de concevoir des interventions adaptées pour prévenir l'émergence future d'infections difficiles à traiter.
Volume	4
Issue	4
Page numbers	94-98

References	1. Appelbaum PC. “MRSA—the tip of the iceberg”. <i>Clin Microbiol Infect.</i> 12(Suppl 2):3–10. 2006 2. Brown DF, Edwards DI, Hawkey PM, et al, on behalf of the Joint Working Party of the British Society for Antimicrobial Therapy, Hospital Infection Society, Infection Control Nurses Association. “Guidelines for the laboratory diagnosis and susceptibility testing of methicillin- resistant <i>Staphylococcus aureus</i> (MRSA)”. <i>J Antimicrob Chemother</i> 56:1000–18. 2005 3. Francis JS, Doherty MC, Lopatin U, Johnston CP, Sinha G, Ross T et al. “Severe community-onset pneumonia in healthy adults caused by methicillin-resistant <i>Staphylococcus aureus</i> carrying the Panton-Valentine leukocidin genes”. <i>Clin Infect Dis.</i> 40:100–7. 2005 4. Healy CM, Hulten KG, Palazzi DL, Campbell JR, Baker CJ. “Emergence of new strains of methicillin-resistant <i>Staphylococcus aureus</i> in a neonatal intensive care unit”. <i>Clin Infect Dis.</i> 39:1460–6. 2004 5. Saxen S, Singh K. and Talwar V. “Methicillin Resistance <i>Staphylococcus aureus</i> prevalence in Community in East Delhi area”. <i>Jpn J Infect Dis.</i> 56:54-56. 2003 6. Ojulong J, Mwambu TP, Joloba M, Bwanga F, Kaddu-Mulindwa DH. “Relative prevalence of methicillin resistant <i>Staphylococcus aureus</i> and its susceptibility pattern in Mulago Hospital, Kampala, Uganda”. <i>Tanzan J Health Res</i> 11: 149–153. 2009
------------	--

	7. Moremi N, Mshana SE, Kamugisha E, Kataraihya J, Tappe D, et al. “Predominance of methicillin resistant <i>Staphylococcus aureus</i> -ST88 and new ST1797 causing wound infection and abscesses”. <i>J Infect Dev Ctries</i> 6: 620–625. 2012 8. Ojikan-Odeke. M.Med. dissertation on “Hand infections in Uganda”. 19-40. 1978 9. Mugisa D.B. M. Med Surgery dissertation on “Complications following laparotomy in Mulago Hospital”. 59-156. 1988 10. Olaro .C. M.Med Surgery dissertation on “Study to assess risk factors for Postoperative Complication following abdominal Surgery in Mulago hospital”. 1-3. 1999 11. Kitara LD, Anywar AD, Acullu D, Odongo-Aginya E, Aloyo J, Fendu M. “Antibiotic susceptibility of <i>Staphylococcus aureus</i> in suppurative lesions in Lacor Hospital, Uganda”. <i>African Health Sciences</i>. 11 (S1): S34 - S39. 2011 12. Shanson DC. “Antibiotic Resistant <i>Staph aureus</i>”. <i>J Hospit infect</i>. 2:11-16. 1981 13. Boyce JM. “Increasing prevalence of MRSA in the USA”. <i>J Infect Control and Hospital Epidemiol</i>. 11:639-642. 1990 14. Kateete D, Kimani C, Katabazi F, Okeng A, Okee M, Nanteza A, Joloba M, Najjuka F. “Identification of <i>Staphylococcus aureus</i>: DNase and Mannitol salt agar improve the efficiency of the tube coagulase
--	--

	test”. <i>Annals of Clinical Microbiology and Antimicrobials</i> 9(1):23. 2010 15. Bull AL, Worth LJ, Richards MJ. “Impact of Vancomycin Surgical Antibiotic Prophylaxis on the Development of Methicillin-Sensitive Staphylococcus aureus Surgical Site Infections: Report From Australian Surveillance Data (VICNISS)”. <i>Annals of surgery</i>. 256 (6): 1089-1092. 2012 16. Kateete DP, Namazzi S, Okee M, Okeng A, Baluku H, Musisi NL, et al. “High prevalence of methicillin resistant Staphylococcus aureus in the surgical units of Mulago hospital in Kampala, Uganda”. <i>BMC Research Notes</i>. 4:326. 2011 17. J Ojulong, T Mwambu, M Jolobo, E Agwu, F Bwanga, C Najjuka, D Kaddu-Mulindwa. “Prevalence of Methicillin resistant Staphylococcus aureus (MRSA) among isolates from surgical site infections in Mulago hospital- Kampala, Uganda”. <i>The Internet Journal of Infectious Diseases</i>. Volume 7 Number 2. 2008 18. Jean-Marie Liesse Iyamba, José Mulwahali Wambale, Cyprien Mbundu Lukukula, Ntongo za Balega Takaisi-Kikuni. “High prevalence of methicillin resistant staphylococci strains isolated from surgical site infections in Kinshasa”. <i>Pan African Medical Journal</i>. 18:322. 2014 19. Bashir Mwambi, Jacob Iramiot, Freddie Bwanga, Marthae Nakaye, Herbert Itabangi and Joel Bazira. “Clindamycin Resistance among
--	--

	Staphylococcus Aureus Isolated at Mbarara Regional Referral Hospital, in South Western Uganda”. <i>British Microbiology Research Journal</i> 4(12): 1335-1344. 2014 20. Jeremiah Seni, Christine F Najjuka, David P Kateete, Patson Makobore, Moses L Joloba, Henry Kajumbula, <i>et al.</i> “Antimicrobial resistance in hospitalized surgical patients: a silently emerging public health concern in Uganda”. <i>BMC Research Notes</i> 6:298. 2013 21. Najat Buzaid, Abdel-Naser Elzouki, Ibrahim Taher, Khalifa Sifaw Ghenghesh. “Methicillin-resistant Staphylococcus aureus (MRSA) in a tertiary surgical and trauma hospital in Benghazi, Libya”. <i>J Infect Dev Ctries.</i> 5(10):723-726. 2011 22. Ching Jou Lim, Allen C. Cheng, Jacqueline Kennon, Denis Spelman, Dayna Hale, Gabrielle Melican <i>et al.</i> “Prevalence of multidrug-resistant organisms and risk factors for carriage in long-term care facilities: a nested case–control study”. <i>J Antimicrob Chemother</i> 2014. 23. Meng Xiao, He Wang, Ying Zhao, Lei-Li Mao, Mitchell Brown, Yun-Song Yu <i>et al.</i> “National Surveillance of Methicillin-Resistant <i>Staphylococcus aureus</i> in China Highlights a Still-Evolving Epidemiology with 15 Novel Emerging Multilocus Sequence Types”. <i>Journal of Clinical Microbiology</i> 51 (11): 3638–3644. 2013 24. Alain C. Juayang, Gemma B. de los Reyes, April Joy G. de la Rama, Christine T. Gallega. “Antibiotic Resistance Profiling of <i>Staphylococcus aureus</i> Isolated from Clinical Specimens in a Tertiary
--	--

	Hospital from 2010 to 2012”. <i>Interdisciplinary Perspectives on Infectious Diseases</i> 898457: 4. 2014 25. Obianuju Onelum, Babatunde Odetoyin, Anthony Onipede and Adesola Oyelese. “The Role of Methicillin-Resistant <i>Staphylococcus aureus</i> in Clinical infections in Obafemi Awolowo University Teaching Hospitals Complex, Ile-Ife, South Western Nigeria”. <i>Journal of Microbiology & Experimentation</i>. 2(2): 00041. 2015 26. Rita Khanal, Prakash Sah, Pramila Lamichhane, Apsana Lamsal, Sweety Upadhaya and Vijay Kumar Pahwa. “Nasal carriage of methicillin resistant <i>Staphylococcus aureus</i> among health care workers at a tertiary care hospital in Western Nepal”. <i>Antimicrobial Resistance and Infection Control</i>. 4:39. 2015 27. Cosgrove SE, Qi Y, Kaye KS, Harbarth S, Karchmer AW, Carmeli Y. “The impact of methicillin resistance in <i>Staphylococcus aureus</i> bacteremia on patient outcomes: mortality, length of stay, and hospital charges”. <i>Infect Control Hosp Epidemiol</i>. 26:166–74. 2005 28. Gastmeier P, Schwab F, Sohr D, Behnke M, Geffers C. “Reproducibility of the surveillance effect to decrease nosocomial infection rates”. <i>Infect Control Hosp Epidemiol</i>. 30:993–9. 2009 29. G. Sganga, C. Tascini, E. Sozio, M. Carlini, P. Chirletti, F. Cortese et al. “Focus on the prophylaxis, epidemiology and therapy of methicillin-resistant <i>Staphylococcus aureus</i> surgical site infections and
--	--

	a position paper on associated risk factors: the perspective of an Italian group of surgeons”. <i>World Journal of Emergency Surgery</i> 11:26. 2016 30. Rubin RJ, Harrington CA, Poon A, Dietrich K, Greene JA, Moiduddin A. “The economic impact of Staphylococcus aureus infection in New York City hospitals”. <i>Emerg Infect Dis.</i> 5:9–17. 1999 31. Pandya N, Chaudhary A, Mehta S, Parmar R. “Characterization of Methicillin Resistant Staphylococcus aureus from various clinical samples at Tertiary care hospital of rural Gujarat”. <i>J Res Med Den Sci.</i> 2(3):49-53. 2014
--	--